

MRCLEAN-R: INTERVENTION

CASE REGISTRATION FORM



Date: ___ / ___ / _____ (dd/mm/yyyy)

1st interventionalist:

2nd interventionalist:

3rd interventionalist:

Paste patient sticker or write down:

Name:
Gender:
Patient #:
Birth date:

1. Intervention

- Catheterization only
- DSA only
- Intervention

If no intervention is performed, please explain:

.....
.....

2. eTICI-score before intra-arterial treatment:

- 0: No perfusion or antegrade flow beyond site of occlusion
- 1: Penetration but not perfusion. Contrast penetration exists past the initial obstruction but with minimal filling of the normal territory
- 2A: Some perfusion with distal branch filling of <50% of territory vascularized
- 2B: Substantial perfusion with distal branch filling of ≥50% of territory visualized
- 2C: Near-complete perfusion except for slow flow in a few distal cortical vessels or presence of small distal cortical emboli
- 3: Complete perfusion with normal filling of all distal branches

3. Location of occlusion:

- M1 M2 A1 A2 ICA ICA-T VA BA P1 other:
- Side: Left Right

4a. General anesthesia during procedure:

- no yes

4b. Conscious sedation during procedure:

- no yes

5. Medication administered during procedure:

- heparin Total dose: IU
- abciximab (Reopro) Total dose: mg
- other Total dose:

6. Treatment with mechanical device:

- Guiding catheter: 6F 8F Balloon: no yes

-
- Stent-retriever: Solitaire attempts:
 - Revive attempts:
 - Catch attempts:
 - TREVO attempts:
 - Other: attempts:

Aspiration device: attempts:

MERCI-retriever attempts:

Other: attempts:

Please specify which stent was used first:

ATTENTION: TURN OVER! Please fill out backside as well!

7. Treatment with **intra-arterial** thrombolytic: no urokinase, dose: alteplase, dose:.....

8. Time

- Arrival patient in angio-suite: (hh:mm)

- Groin puncture:

- First attempt with device:

- Time of first recanalization (TICI 2b/3) or last contrast bolus:

- Sheath from groin:

- Total duration of IA treatment (groin puncture to sheath withdrawal) minutes

9. Thrombus extracted for PA: no yes

10. eTICI-score after intra-arterial treatment:
- 0: No perfusion
 - 1: Antegrade reperfusion past the initial occlusion, but limited distal branch filling with little or slow distal reperfusion
 - 2A: Antegrade reperfusion of less than half of the occluded target artery previously ischemic territory (e.g. in 1 major division of the MCA and its territory)
 - 2B: Antegrade reperfusion of more than half of the previously occluded target artery ischemic territory (e.g. in 2 major divisions of the MCA and their territories)
 - 2C: Near-complete perfusion except for slow in a few distal cortical vessels or presence of small distal cortical emboli
 - 3: Complete antegrade reperfusion of the previously occluded target artery ischemic territory, with absence of visualized occlusion in all branches

11. Procedural complications / adverse events (e.g. vasospasm, dissection, perforation or peripheral emboli):

12. Description of procedure (optional):

13. Closure device used:

- angioseal
 - other, please specify type:
- Side: Left Right

14. Have all images been transferred to the PACS system?

Please include baseline & post-intervention AP + lateral imaging up until venous phase.

15. Signature: _____ Date: ___ / ___ / _____ (dd/mm/yyyy)